

Temperature Preconditioning

Susan Lurie¹ and Joshua D. Klein²

¹Department of Postharvest Science and ²Department of Field Crops
Volcani Center, Bet-Dagan, Israel

Introduction: Temperature preconditioning of fruits and vegetables has been practiced for more than 70 yr, since Baker (1939; 1952) described heat treatments for disinfestation of fruit flies in citrus. There is renewed interest in high temperature as a postharvest treatment for control of both insect pests and fungal pathogens in fresh produce. In part, this is because of the deregistration of a number of compounds that have until recently been used for effective control of postharvest disorders. In addition, there is increased consumer demand for produce that has had minimal, or ideally no, chemical treatment.

Heat has fungicidal and well as insecticidal action, but heat regimes that are optimal for insect control may not be optimal for disease control; in some cases they may even be detrimental. A thermal treatment that is developed for fungus or insect control should not damage the commodity being treated.

In fact, in many cases high temperature manipulation before storage may have beneficial effects on the commodity treated. These benefits include slowing the ripening of climacteric fruit and vegetables, enhancing sweetness of produce by increasing the amount of sugars or decreasing acidity, and prevention of storage disorders such as superficial scald on apples and chilling injury on subtropical fruits and vegetables (Lurie, 1998).

Temperature conditioning before storage may also mean an incubation period spent at either ambient temperature of 16 to 25 °C (about 61 to 77 °F) or at a temperature below ambient, but above that which might produce chilling injury, i.e., 5 to 12 °C (about 41 to 54 °F), depending on commodity. This type of temperature manipulation is often referred to as a 'curing' period, and it is used with crops such as potatoes, onions and carrots. Its purpose is generally to enhance resistance of the commodity to pathogen invasion, although it may also enhance resistance to low temperature injury in citrus.

In this chapter we discuss temperature preconditioning treatments according to their purpose, i.e., pathogen, insect or chilling injury control. Most of the methods listed here, however, are still experimental, and have yet to be accepted for routine commercial practice.

Commercial Treatments: The greatest number of temperature manipulations used commercially are based on high temperature treatments (vapor heat or hot forced-air) for insect disinfestation. Temperature regimes are developed specifically for each commodity and insect pest. The accepted procedures for produce entering the U.S. are described in the USDA-APHIS Plant Protection and Quarantine Treatment Manual, which is routinely updated (Animal and Plant Protection Service, 1998). The latest edition of the manual should be consulted for approved treatments for particular commodities or pests.

An example of commercial temperature conditioning for pest control is Mexican-grown mangos, which may be infested with a variety of fruit fly larvae or eggs. Officially authorized treatments are high-temperature, forced-air (HAT) or a hot water dip treatment (HWT) before storage and shipment. In HAT, fruit are heated until their center reaches 48 °C (118 °F). HWT conditions depend on fruit size and vary from 45 to 90 min in water, where the fruit interior reaches 46.1 °C (115 °F).

Vapor heat (VHT) differs from high-temperature, forced-air in that moisture accumulates on the surface of the fruit. The water droplets transfer heat more efficiently than air, allowing the fruit to heat quickly; but there may also be increased physical injury to the fruit. Papayas grown in Hawaii are vapor heat-treated before export to Japan.

Citrus can be disinfested by HAT at 44 °C (111 °F) for 100 min, with an additional 90 min spent raising the temperature to 44°C. The usual disinfestation method, however, is to hold the fruit at low temperature of 0 to 2.2 °C (32 to 36 °F) for 10 to 16 days, before raising the temperature to the normal storage temperature of 6 to 11°C (43 to 52 °F), depending on cultivar. Since citrus is sensitive to chilling,

fruit are generally held at 20 °C (68 °F) or 16 °C (61 °F) for 3 to 5 days before placing at low temperature. This curing treatment decreases fruit susceptibility to chilling injury resulting from the subsequent disinfestation treatment.

Insect Disinfestation: The development and implementation of heat treatments for insect disinfestation have been reviewed thoroughly (Couey, 1989; Paull, 1993). The list below includes treatment regimes that have been reported in the past 10 years (Table 1). More than half the treatments are designed to kill fruit fly eggs or larvae, since their presence requires strict quarantine in most fruit-importing countries. The most recently developed methods include heat treatments in combination with low O₂ or high CO₂ atmospheres.

Antifungal Treatments: Curing is used commercially to increase resistance to pathogen invasion. Potatoes are cured at 12 °C (54 °F) for 10 to 12 days before storage at 4 to 9 °C (39 to 48 °F), depending on cultivar and whether they are designated for industry or home consumption. Sweet potatoes are also cured at 30 °C (86 °F) for 5 days, before storage at 12 °C (54 °F). In both cases the curing period allows for wound healing and deposition of cell wall material to create a physical barrier to pathogens. Kiwifruit also benefit from a curing period. If held at 10 °C (50 °F) before storage at low temperature, they develop fewer rots after storage. Onions can be stored longer if held at 28 °C (82 °F) for 3 days before storage.

The two commercial applications of high temperature antifungal treatments are HWT for papayas (Akamine and Arisumi, 1953), which has been used for almost 50 yr, and a hot water brush treatment that was introduced fairly recently (Fallik, 1996a, 1999; Prusky et al., 1997). The brush system is in use on packing lines for export of corn, mangos, peppers and some citrus from Israel. The machine sprays hot water at 50 to 65 °C (122 to 149 °F) on produce as it moves along on brush rollers. The major benefit appears to be removal of spores and dirt, although hot water combined with brushing also causes surface cracks to be filled in by the natural wax of the commodity, as well as eliciting resistance to pathogens in some cases.

The state of temperature conditioning treatments against fungal pathogens was reviewed by Barkai-Golan and Phillips (1991) and Coates and Johnson (1993). The majority of the regimes listed in Table 2 were developed in the past 5 years. Dips in hot fungicide solution have been used since the 1950s for pathogen control. As various fungicides lose their registration or as pathogens develop resistance, there is increased interest in heat-treating produce in combination with compounds that are generally recognized as safe (GRAS), such as CaCl₂ or sodium carbonate (Table 2).

Physiological Benefits of Conditioning Treatments: Most thermal treatments have been developed as lethal regimes for insects or fungi. Some of these regimes, however, also have prophylactic effects against physiological disorders such as chilling injury (CI). Prevention of CI allows the commodity to be stored longer at lower temperatures, which in turn permits export in ships rather than more costly air-freight. In addition, a pre-shipping heat treatment can allow for low temperature disinfestations of commodities such as citrus, by improving the resistance of fruit to CI generally incurred during this treatment.

Other heat treatments have been developed specifically to maintain postharvest quality, such as increased firmness of apples or decreased yellowing of broccoli, or to protect against other abiotic stresses, such as irradiation disinfestation treatments (Table 3). The physiological mechanisms of these treatments was previously reviewed by Lurie (1998).

Literature Cited:

- Afek, U., J. Orenstein and E. Nuriel. 1999. Steam treatment to prevent carrot decay during storage. *Crop Protection*. 18:639-642.
- Akamine, E.K. and T. Arisumi. 1953. Control of postharvest storage decay of fruits of papaya (*Carica papaya* L.) with special reference to the effect of hot water. *Proc. Amer. Soc. Hort. Sci.* 61:270-274.
- Animal and Plant Health Inspection Service. 1998. Plant protection and quarantine manual. USDA,

APHIS, Hyattsville, MD.

- Armstrong, J., B. Hu and S. Brown. 1995. Single-temperature forced hot-air quarantine treatment to control fruit flies (Diptera: Tephritidae) in papaya. *J. Econ. Entomol.* 88:678-682.
- Baker, A.G. 1939. The basis for treatment of products where fruit flies are involved as a condition for entry into the United States. USDA Circular 551, Washington, DC.
- Baker, A.G. 1952. The vapor-heat process. USDA Yearbook, U.S. Gov. Print. Off., Wash. DC.
- Barkai-Golan, R. and D.J. Phillips. 1991. Postharvest heat treatment of fresh fruits and vegetables for decay control. *Plant Dis.* 75:1085-1089.
- Barkai-Golan, R., R. Padova, I. Ross, M. Lapidot, H. Davidson and A. Copel. 1993. Combined hot water and radiation treatments to control decay of tomato fruits. *Sci. Hort.* 56:101-105.
- Ben-Yehoshua, S., B. Shapiro and R. Moran. 1987. Individual seal-packaging enables the use of curing at high temperatures to reduce decay and heat injury of citrus fruits. *HortScience* 22:777-783.
- Chan, H.T. and E. Linse. 1989. Conditioning cucumbers for quarantine heat treatments. *HortScience* 24:985-989.
- Chervin, C., S. Kulkarni, S. Kreidl, F. Birrell and D. Glenn. 1997. A high temperature/low oxygen pulse improves cold storage disinfestation. *Postharv. Biol. Technol.* 10:239-245.
- Coates, L.M. and G.I. Johnson. 1993. Effective disease control in heat-disinfested fruit. *Postharv. News Info.* 4:35N-40N.
- Coates, L.M., G. I. Johnson and A. Cooke. 1993. Postharvest disease control in mangoes using high humidity hot air and fungicide treatments. *Ann. Appl. Biol.* 123:441-448.
- Couey, H.M. 1989. Heat treatment for control of postharvest diseases and insect pests of fruits. *HortScience.* 24:198-201.
- Dentener, P.R., S. M. Alexander, P.J. Lester, R.J. Petry, J.H. Maindonald and R.M. McDonald. 1996. Hot air treatment for disinfestation of lightbrown apple moth and longtailed mealy bug on persimmons. *Postharv. Biol. Technol.* 8:143-152.
- Dentener, P.R., K.V. Bennett, L.E. Hoy, S. E. Lewthwaite, P.J. Lester, J.H. Maindonald and P.G. Connolly. 1997. Postharvest disinfestation of lightbrown apple moth and longtailed mealybug on persimmons using heat and cold. *Postharv. Biol. Technol.* 12:255-264.
- Dentener, P. R., S.E. Lewthwaite, J.H. Maindonald and P.G. Connolly. 1998. Mortality of twospotted spider mite (Acari: Tetranychidae) after exposure to ethanol at elevated temperatures. *J. Econ. Entomol.* 91:767-772.
- Dentener, P.R., S.E. Lewthwaite, K.V. Bennett, J.H. Maindonald and P.G. Connolly. 2000. Effect of temperature and treatment conditions on the mortality of *Epiphyas postvittana* (Lepidoptera: Tortricidae) exposed to ethanol. *J. Econ. Entomol.* 93:519-525.
- Fallik, E., J. D. Klein, S. Grinberg, E. Lomaniec, S. Lurie and A. Lalazar. 1993. Effect of postharvest heat treatment of tomatoes on fruit ripening and decay caused by *Botrytis cinerea*. *Plant Dis.* 77:985-988.
- Fallik, E., Y. Aharoni, O. Yekutieli, A. Wiseblum, R. Regev, H. Beres and E. Bar Lev. 1996a. A method for simultaneously cleaning and disinfecting agricultural produce. Israel Patent Application No. 116965
- Fallik, E., S. Grinberg, S. Alkalai and S. Lurie. 1996b. The effectiveness of postharvest hot water dipping on the control of gray and black molds in sweet red pepper (*Capsicum annuum*). *Plant Pathol.* 45:644-649.
- Fallik, E., S. Grinberg, M. Gambourg, J.D. Klein and S. Lurie. 1996c. Prestorage heat treatment reduces pathogenicity of *Penicillium expansum* in apple fruit. *Plant Pathol.* 45:92-97.
- Fallik, E., S. Grinberg, S. Alkalai, O. Yekutieli, A. Weisblum, R. Regev, H. Beres and E. Bar-Lev. 1999. A unique rapid hot water treatment to improve storage quality of sweet pepper. *Postharv. Biol. Technol.* 15:25-32.
- Florissen, P., J.S. Ekman, C. Blumenthal, W.B. McGlasson, J. Conroy and P. Holford. 1996. The effects of short heat treatments on the induction of chilling injury in avocado fruit (*Persea americana* Mill). *Postharv. Biol. Technol.* 8:129-141.

- Follet, P.A. and Z. Gabbard. 1999. Efficacy of the papaya vapor heat quarantine treatment against white peach scale in Hawaii. *HortTechnology* 9:506.
- Forney, C.F. 1995. Hot water dips extend shelf-life of fresh broccoli. *HortScience* 30:1054-1057.
- Garcia, J.M., C. Aguilera and A.M. Jimenez. 1996. Grey mold in and quality of strawberry fruit following postharvest heat treatment. *HortScience* 31:255-257.
- Gonzalez-Aguilar, G.A., L. Zacarias and M.T. Lafuente. 1998. Ripening affects high-temperature-induced polyamines and their changes during cold storage of hybrid Fortune mandarins. *J. Agric. Fd Chem.* 46:3503-3508.
- Hallman, G.J., J.J. Gaffney and J.L. Sharp. 1990. Vapor heat treatment for grapefruit infested with Caribbean fruit fly (Diptera: Tephritidae). *J. Econ. Entomol.* 83:1475-1478.
- Heather, N.W., R.J. Corcoran and R.A. Kopittke. 1997. Hot air disinfestation of Australian 'Kensington' mangoes against two fruit flies (Diptera: Tephritidae). *Postharv. Biol. Technol.* 10:99-105.
- Hong, G., G. Peiser and M.I. Cantwell. 2000. Use of controlled atmospheres and heat treatment to maintain quality of intact and minimally processed green onions. *Postharv. Biol. Technol.* 20:53-61.
- Hoy, L.E. and D. C. Whiting. 1998. Mortality responses of three leafroller (Lepidoptera: Tortricidae) species on kiwifruit to a high-temperature controlled atmosphere treatment. *N.Z. J. Crop Hort. Sci.* 26:11-15.
- Jacobi, K.K., Wong, L.S. and J.E. Giles. 1993. Lychee (*Lichi chinensis* Sonn.) fruit quality following vapour heat treatment and cool storage. *Postharv. Biol. Technol.* 3:111-119.
- Jacobi, K.K., J. Giles, E. MacRae and T. Wegrzyn. 1995a. Conditioning 'Kensington' mango with hot air alleviates hot water disinfestation injuries. *HortScience* 30:562-565.
- Jacobi, K.K., L.S. Wong and J.E. Giles. 1996. Postharvest quality of zucchini (*Cucurbita pepo* L.) following high humidity hot air disinfestation treatments and cool storage. *Postharv. Biol. Technol.* 7:309-316.
- Jacobi, K.K. and J.E. Giles. 1997. Quality of 'Kensington' mango (*Mangifera indica* Linn.) fruit following combined vapor heat disinfestation and hot water disease control. *Postharv. Biol. Technol.* 12:285-292.
- Jang, E.B. 1996. Systems approach to quarantine security: postharvest application of sequential mortality in the Hawaiian grown 'Sharwil' avocado system. *J. Econ. Entomol.* 89:950-956.
- Jessup, A.J. 1994. Quarantine disinfestation of 'Hass' avocados against *Bactrocera tryoni* (Diptera: Tephritidae) with a hot fungicide dip followed by cold storage. *J. Econ. Entomol.* 87:127-130.
- Kim, J.J., S. Ben-Yehoshua, B. Shapiro, Y. Henis and S. Carmeli. 1991. Accumulation of scoparone in heat-treated lemon fruit inoculated with *Penicillium digitatum* Sacc. *Plant Physiol.* 97:880-885.
- Klein, J.D. and S. Lurie. 1992. Prestorage heating of apple fruit for enhanced postharvest quality: interaction of time and temperature. *HortScience* 27:326-328.
- Klein, J., W. Conway, B. Whitaker and C. Sams. 1997. *Botrytis cinerea* decay in apples is inhibited by postharvest heat and calcium treatments. *J. Amer. Soc. Hort. Sci.* 122:91-94.
- Lay-Yee, M. and D.C. Whiting. 1996. Response of 'Hayward' kiwifruit to high-temperature controlled atmosphere treatments for control of two-spotted spider mite (*Tetranychus urticae*). *Postharv. Biol. Technol.* 7:73-81.
- Lay-Yee, M., S. Ball, S.K. Forbes and A.B. Woolf. 1997a. Hot water treatment for insect disinfestation and reduction of chilling sensitivity of 'Fuyu' persimmon. *Postharv. Biol. Technol.* 10:81-89.
- Lay-Yee, M., D.C. Whiting and K.J. Rose. 1997b. Response of 'Royal Gala' and 'Granny Smith' apples to high-temperature controlled atmosphere treatments for control of *Epiphyas postvittana* and *Nysius huttoni*. *Postharv. Biol. Technol.* 12:127-136.
- Lester, P.L., P.R. Dentener, R.J. Petry and S.M. Alexander. 1995. Hot-water immersion for disinfestation of lightbrown apple moth (*Epiphyas postvittana*) and longtailed mealy bug (*Pseudococcus longispinus*) on persimmons. *Postharv. Biol. Technol.* 6:349-356.
- Lester, P.J., P.R. Dentener, K.V. Bennett and P.G. Connolly. 1997. Postharvest disinfestations of diapausing and non-diapausing twospotted spider mite (*Tetranychus urticae*) on persimmons: hot

- water immersion and coolstorage. *Entom. Exper. Appl.* 83:189-193.
- Lurie, S. 1998. Postharvest heat treatments. *Postharv. Biol. Technol.* 14: 257-269.
- Lurie, S. and J.D. Klein. 1991. Acquisition of low temperature tolerance in tomatoes by exposure to high temperature stress. *J. Amer. Soc. Hort. Sci.* 116:1007-1012.
- Lurie, S., J.D. Klein and R. Ben-Arie. 1990. Postharvest heat treatment as a possible means of reducing superficial scald of apples. *J. Hort. Sci.* 65:503-509.
- Lurie, S. M. Laamim, Z. Lapsker and E. Fallik. 1997. Heat treatments to decrease chilling injury in tomato fruit. Effects on lipids, pericarp lesions and fungal growth. *Physiol. Plant.* 100:297-302.
- McCollum, T.G., S. D'Aquino and R.E. McDonald. 1993. Heat treatment inhibits mango chilling injury. *HortScience* 28:197-198.
- McCollum, G., H. Doostdar, R. Mayer and R. McDonald. 1995. Immersion of cucumber in heated water alters chilling-induced physiological changes. *Postharv. Biol. Technol.* 6:55-64.
- McDonald, R.E., T.G. McCollum and E.A. Baldwin. 1998. Heat treatment of mature-green tomatoes: differential effects of ethylene and partial ripening. *J. Amer. Soc. Hort. Sci.* 123:457-462.
- McDonald, R.E., T.G. McCollum and E.A. Baldwin. 1999. Temperature of water heat treatments influences tomato fruit quality following low-temperature storage. *Postharv. Biol. Technol.* 16:147-155.
- McGuire, R.G. 1991. Concomitant decay reductions when mangoes are treated with heat to control infestations of Caribbean fruit fly. *Plant Dis.* 75:946-949.
- McGuire, R.G. 1997. Market quality of guavas after hot-water quarantine treatment and application of carnauba wax coating. *HortScience* 32:271-274.
- McLaren, G.F., R.M. McDonald, J.A. Fraser, R.R. Marshall, K.J. Rose and A.J. Ford. 1997. Disinfestation of New Zealand flower thrips from stonefruit using hot water. *Acta Hort.* 464:524.
- Mencarelli, F., B. Ceccantoni, A. Bolini and G. Anelli. 1993. Influence of heat treatment on the physiological response of sweet pepper kept at chilling temperature. *Acta Hort.* 343:238-243.
- Miller, W.R., R.E. McDonald and J.L. Sharp. 1991. Quality changes during storage and ripening of 'Tommy Atkins' mangos treated with heated forced air. *HortScience* 26:395-397.
- Neven, L. and E. Mitcham. 1996. CATTS: controlled atmosphere/temperature treatment system. A novel tool for the development of quarantine treatments. *Amer. Entomol.* 42:56-59.
- Neven, L.G. and S.R. Drake. 2000. Comparison of alternative postharvest quarantine treatments for sweet cherries. *Postharv. Biol. Technol.* 20:107-114.
- Neven, L.G., L.M. Rehfield and K. C. Shellie. 1996. Moist and vapor forced air treatments of apples and pears: effects on the mortality of fifth instar codling moth (Lepidoptera: Tortricidae). *J. Econ. Entomol.* 89:700-704.
- Nishijima, K.A., K. Miura, J.W. Armstrong, S.A. Brown, B.K.S. Hu. 1992. Effect of forced, hot-air treatment of papaya fruit on fruit quality and incidence of postharvest diseases. *Plant Dis.* 76:723-727.
- Nishijima, K., H. Chan, S. Sanxter and E. Linse. 1995. Reduced heat shock period of 'Sharwil' avocado for cold tolerance in quarantine cold treatment. *HortScience* 30:1052-1053.
- Paull, R.E. 1994. Response of tropical horticultural commodities to insect disinfestation treatments. *HortScience* 29:988-996.
- Paull, R.E. and N.J. Chen. 1999. Heat treatment prevents postharvest geotropic curvature of asparagus spears (*Asparagus officinallis* L.). *Postharv. Biol. Technol.* 16:37-41.
- Porat, R., D. Pavoncello, J. Peretz, S. Ben-Yehoshua and S. Lurie 1999. Effects of various heat treatments on the induction of cold tolerance and on the postharvest qualities of 'Star Ruby' grapefruit. *J. Amer. Soc. Hort. Sci.* 124:184-188.
- Porat, R., A. Daus, B. Weiss, L. Cohen, E. Fallik, and S. Droby. 2000. Reduction of postharvest decay in organic citrus fruit by a short hot water brushing treatment. *Postharv. Biol. Technol.* 18:151-157.
- Prusky, D., Y. Fuchs, I. Kobiler, I. Roth, A. Weksler, Y. Shalom, E. Fallik, G. Zauberman, E. Pesis, M. Akerman, O. Ykutiely, A. Weisblum, R. Regev and L. Artes. 1999. Effect of hot water brushing,

- prochloraz treatment and waxing on incidence of black spot decay caused by *Alternaria alternata* in mango fruits. *Postharv. Biol. Technol.* 15:165-174.
- Ranagann, B., G.S.V. Raghavan and A.C. Kushalappa. 1998. Hot water dipping to enhance storability of potatoes. *Postharv. Biol. Technol.* 13:215-223.
- Reyes, M.E.Q., W. Nishijima and R.E. Paull. 1998. Control of crown rot in 'Santa Catarina Prata' and 'Williams' banana with hot water treatment. *Postharvest Biol. Technol.* 14:71-75.
- Rodov, V., S. Ben-Yehoshua, R. Albagli and D.Q. Fang. 1995. Reducing chilling injury and decay of stored citrus fruit by hot water dips. *Postharv. Biol. Technol.* 5:119-127.
- Sams, C.E., W.S. Conway, J.A. Abbott, R.J. Lewis and N. Ben-Shalom. 1993. Firmness and decay of apples following postharvest pressure infiltration of calcium and heat treatment. *J. Amer. Soc. Hort. Sci.* 118:623-627.
- Schirra, M. and G. D'Hallewin. 1997. Storage performance of Fortune mandarins following hot water dips. *Postharv. Biol. Technol.* 10:229-238.
- Schirra, M., G. Barbera, S. D'Aquino, T. La Mantia, and R.E. McDonald. 1996. Hot dips and high-temperature conditioning to improve shelf quality of late-crop cactus pear fruit. *Trop. Sci.* 36:159-165.
- Schirra, M., M. Agabbio, S. d'Aquino and T.G. McCollum. 1997. Postharvest heat conditioning effects on early ripening 'Gialla' cactus pear fruit. *HortScience* 32:702-704.
- Schirra, M., P. Cabras, A. Angioni, G. D'hallewin, R. Ruggiu and E.V. Minelli. 1997. Effect of heated solutions on decay control and residues of imazalil in lemons. *J. Agric. Fd Chem.* 45:4127-4130.
- Sharp, J.L. and W.P. Gould. 1994. Control of Caribbean fruit fly (Diptera: Tephritidae) in grapefruit by forced hot air and hydrocooling. *J. Econ. Entomol.* 87:131-133.
- Sharp, J.L. and R.G. McGuire. 1996. Control of Caribbean fruit fly (Diptera: Tephritidae) in navel orange by forced hot air. *J. Econ. Entomol.* 89:1181-1185.
- Shellie, K.C., R.L. Mangan and S.J. Ingle. 1997. Tolerance of grapefruit and Mexican fruit fly larvae to heated controlled atmospheres. *Postharv. Biol. Technol.* 10:179-186.
- Shellie, K.C. 1998. Reduction of green mold on grapefruit after hot forced-air quarantine treatment. *Plant Dis.* 82:380-382.
- Smilanick, J.L., D.A. Margosan and D.J. Henson. 1995. Evaluation of heated solutions of sulfur dioxide, ethanol, and hydrogen peroxide to control postharvest green mold of lemons. *Plant Dis.* 79:742-747.
- Smilanick, J.L., B.E. Mackey, R. Reese, J. Usall and D.A. Margosan. 1997. Influence of concentration of soda ash, temperature, and immersion period on the control of postharvest green mold of oranges. *Plant Dis.* 80:230-234.
- Sonderstrom, E.L., D.G. Brandl and B.E. Mackay. 1993. High temperature for control of *Asynonychus godmani* (Coleoptera: Curculionidae) eggs on lemon fruit. *J. Econ. Entomol.* 86:1773-1780.
- Tian, M.S., A.B. Woolf, J.H. Bowen and I.B. Ferguson. 1996. Changes in color and chlorophyll fluorescence of broccoli florets following hot water treatment. *J. Amer. Soc. Hort. Sci.* 121:310-313.
- Tian, M.S., T. Islam, D.G. Stevenson and D.E. Irving. 1997. Color, ethylene production, respiration, and compositional changes in broccoli dipped in hot water. *J. Amer. Soc. Hort. Sci.* 122:112-116.
- Wang, C.Y. 1994. Combined treatment of heat shock and low temperature conditioning reduces chilling injury in zucchini squash. *Postharv. Biol. Technol.* 4:65-73.
- Wang, C.Y. 1998. Heat treatment affects postharvest quality of kale and collard, but not of Brussels sprouts. *HortScience* 33:881-883.
- Whiting, D.C. and L.E. Hoy. 1997. High temperature controlled atmosphere and air treatments to control obscure mealybug (Hemiptera: Pseudococcidae) on apples. *J. Econ. Entomol.* 90:546-550.
- Whiting, D.C., L.E. Jamieson, K.J. Spooner and M. Lay-Yee. 1999. Combination of high-temperature controlled atmosphere and cold storage as a quarantine treatment against *Ctenopseustis obliquana* and *Epiphyas postvittana* on 'Royal Gala' apples. *Postharv. Biol. Technol.* 16:119-126.
- Woolf, A.B., C.B. Watkins, J.H. Bowen, M. Lay-Yee, J.H. Maindonald and I.B. Ferguson. 1995. Reducing external chilling injury in stored 'Hass' avocados with dry heat treatments. *J. Amer. Soc.*

Hort. Sci. 120:1050-1056.

Woolf, A.B. and M. Lay-Yee. 1997. Pretreatments at 38 °C of 'Hass' avocado confer thermotolerance to 50 °C hot water treatments. HortScience 32:705-708.

Woolf, A.B., S. Ball, K. Spooner, M. Lay-Yee, I.B. Ferguson, C.B. Watkins, A. Gunson and S. Forbes. 1997. Reduction of chilling injury in the sweet persimmon 'Fuyu' during storage by dry air heat treatments. Postharv. Biol. Technol. 11:155-164.

Table 1.**Insect
Fruit flies**

Latin name	Fruit	Regime	Temperature/Time	Reference
<i>Anastrepha suspensa</i>	grapefruit mango orange	HAT* HAT HAT	51.5 °/125 min	Sharp & Gould '94 Miller et al. '91 Sharp & McGuire '96
<i>Ceratitis capitata</i>	avocado mango papaya	HAT VHT HAT	40 °/24 h 47 °/15 min 47.2 ° at pulp for 3.5 h	Jang '96 Heather et al. '97 Armstrong et al. '95
<i>Dacus cucurbitae</i> <i>Bactrocera cucurbitae</i>	avocado cucumber papaya zucchini	HAT HAT then HWT HAT VHT	40 °/24 h 32.5 °/24 h then 45-46 °/50-60 min 47.2 ° at pulp for 3.5 h	Jang '96 Chan & Linse '89 Armstrong et al. '95 Jacobi et al. '96
<i>Anastrepha ludens</i> <i>Bactrocera cucumis</i>	grapefruit zucchini	HAT & CA VHT	44 °/2 h in 1% O ₂ 45 °/30 min	Shellie et al. '97 Jacobi et al. '96
<i>Dacus dorsalis</i> <i>Bactrocera dorsalis</i>	cucumber papaya	HAT then HWT HAT	32.5 °/24 h then 45-46 °/50-60 min 47.2 ° at pulp for 3.5 h	Chan & Linse '89 Armstrong et al. '95
<i>Bactrocera payapae</i>	mango	VHT	47 °/15 min	Heather et al '97
<i>Bactrocera tyroni</i>	avocado litchi mango	HWT & benomyl VHT VHT HWT then VHT	46 °C/3 min then 1 °/7 days 45 °/30 min 46.5 °/10 min 53 °C/15 min then 47°C/15 min	Jessup '94 Jacobi et al. '93 Heather et al. '97 Jacobi et al. '95 Jacobi & Giles '97

Other Insects

<i>Cydia pomonella</i>	apple cherry pear	HAT or VHT HAT & CA HAT or VHT HAT & CA	44 °/120 min then 0 °/4 weeks 47 °/44 min in 1% O ₂ ; 15% CO ₂ 44 °/120 min then 0°/4 weeks 30 °/ 30 h in 0.3% O ₂	Neven et al. '96 Neven & Mitcham '96 Neven & Drake '00 Neven et al. '96 Chervin et al. '97
<i>Asynonychus godmani</i>	lemon	HWT	52 °/8 min	Soderstrom et al. '93
<i>Cnephasia jactatana</i> <i>Ctenopseustis obliquana</i>	apple kiwifruit	HAT & CA HAT & CA	40 °/10 h in 0.4% O ₂ 45 °/5 h in 0.4% O ₂ 40 °/5-7 h in 0.4% O ₂ 40 °/6 h in 2% O ₂ ; 5% CO ₂	Whiting et al. '99 Whiting et al '97 Hoy & Whiting '98
<i>Epiphyas postvittana</i>	apple kiwifruit pear	HAT & CA HWT & ethanol HAT & CA HAT & CA	40 °/17-20 h in 1.2% O ₂ ; 1% CO ₂ 45 °/13 min in 50% ethanol 30 °/30 h in 0.3% O ₂	Lay-Yee et al. '97 Dentener et al. '00 Hoy & Whiting '98 Chervin et al. '97
<i>Pseudococcus longispinus</i>	persimmon	HWT HAT	48 °/26 min or 50 °/22 min	Lester et al. '95 Dentener et al. '96,'97
<i>Thrips obscuratus</i>	apricot nectarine peach	HWT	48 °/3 min then 50 °/2 min	McLaren et al. '97
<i>Pseudococcus affinis</i>	apple	HAT & CA	40 °/10 h in 0.4% O ₂ 45 °/5 h in 0.4% O ₂	Whiting & Hoy '97
<i>Grapholita molesta</i>	pear	HAT & CA	30 °/30 h in 0.3% O ₂	Chervin et al. '97
<i>Tetranychus urticae</i>	apples kiwifruit perimmon	HWT & ethanol HAT & CA HWT	45 °/13 min in 50% ethanol 44 °/211 min 47 °/67 min	Dentener et al. '98 Lay-Yee & Whiting '96 Lester et al. '97
<i>Pseudaulacaspis pentagona</i>	papaya	VHT	47.2 °/4 h	Follet & Gabbard '99

Table 2. Thermal treatment of horticultural commodities for eradication of and protection from fungal pathogens.

Fungus	Common name	Crop	Regime	Temperature/Time	Reference
<i>Alternaria alternata</i>	Black spot	carrot	HWB	100 °/3 sec	Afek et al. '99
		mango	HWB	60-70 °/15-20 sec	Prusky et al. '99
	Black mold	pepper	HWT	50 °/3 min	Fallik et al. '96b
<i>Botrytis cinerea</i>	Grey mold	Apple	HAT & CaCl ₂	38 °/4 days and CaCl ₂ dip	Klein et al. '97
		pepper	HWT	50 °/3 min	Fallik et al. '96b
		strawberry	HWT	45 °/15 min	García et al. '96
		tomato	HWT	50 °/2 min	Barkai-Golan et al. '93
			HAT	38 °/2 days	Fallik et al. '93
<i>Botryodiplodia theobromae</i>	Stem and surface rots	papaya	HAT	49 °/20 min 32 °/30 min then 49 °/20 min	Nishijima et al. '92
<i>Chalara paradoxa</i>	Crown rot	banana	HWT	45 °/20 min or 50 °/10 min	Reyes et al. '98
<i>Colletotrichum gloeosporioides</i>	Anthracnose	mango	VHT	46-48 °/24 sec - 8 min	Coates et al. '93
			HWT HAT	51.5 °/125 min	McGuire '91 Miller et al. '91
<i>Diplodia natalensis</i>	Stem end rot	mango	HAT HWT	51.5 °/125 min	Miller et al. '91 McGuire '91
<i>Mycospharella spp.</i>	Stem and surface rots	papaya	HAT	49 °/20 min 42 °/30 min then 49°/20 min	Nishijima et al. '92
<i>Penicillium digitatum</i>	Green mold	grapefruit	HAT HWB	46 °/6 h 59-62 °/15 sec	Shellie '98 Porat et al. '00
		lemon	HAT HWT & Na ₂ CO ₃	36 °/3 days 45 °/150 sec + 2% Na ₂ CO ₃	Kim et al. '91 Smilanick et al. '97
		orange	HWT	53 °/3 min	Schirra et al. '97
			HWT & Na ₂ CO ₃	41-43 °/1-2 min + 6% Na ₂ CO ₃	Smilanick et al. '97
<i>Penicillium expansum</i>	Blue mold	Apple	HAT & CaCl ₂ HAT	38 °/4 days + 4% CaCl ₂ 38°/4 days	Sams et al. '93 Fallik et al. '96c
<i>Penicillium italicum</i>	Blue mold	cactus pear	HAT or HWT	38 °/24 h or 55 °/5 min	Schirra et al. '96
<i>Penicillium spp.</i>		lemon	HWT & imazalil	50 °/3 min + imazalil	Schirra et al. '97
<i>Rhizopus stolonifer</i>		tomato	HWT	50 °/2 min	Barkai-Golan et al. '93

Table 3. Physiological benefits of thermal treatments for horticultural crops.

1. Chilling injury

Crop	Phenomenon/Appearance	Regime	Temperature/Time	Reference
Apple	scald	HAT*	38 °/4 days or 42 °/2 days	Lurie et al. '90
Avocado	skin browning internal browning, pitting	HAT then	38 °/3-10 h then	Woolf et al. '95
		HWT	40 °/30 min	Florissen et al. '96
		HWT	38 °/60 min	Woolf et al. '97
Cactus pear	rind pitting, brown staining	HAT or HWT	38 °/24 h or 55 °/5 min	Schirra et al. '96
Citrus	rind pitting	HAT	34-36 °/48-72 h	Ben -Yehoshua et al. '87 Gonzalez-Aguilare et al. '98
		HWT	50-54 °/3 min 53 °/2-3 min	Schirra & D'hallewin '97 Rodov et al. '95
		HWB	59-62 °/15-30 sec	Porat et al. '99
Mango	pitting	HAT	38 °/2 days 54 °/20 min	McCullum et al. '93 Jacobi et al. '95
Persimmon	gel formation	HWT	47 °/90-120 min; 50 °/30-45 min; 52 °/20-30 min	Lay-Yee et al. '97
		HAT		Woolf et al. '97
Green pepper	pitting	HAT	40 °/20 h	Mencarelli et al. '93
Cucumber	pitting	HWT	42 °/30 min	McCullum et al. '95
Tomato	pitting	HAT	38 °/2-3 days	Lurie & Klein '91
		HWT	48 °/2 min	Lurie et al. '97
			42 °/60 min	McDonald et al. '98, '99
Zucchini	pitting	HWT	42 °/30 min	Wang '94

2. Improved postharvest quality

Commodity	Parameter/attribute	Regime	Temperature/Time	Reference
Apple	increased firmness	HAT	38 °4 days; 42 °/2 days	Klein & Lurie '92
Asparagus	inhibited curvature	HWT	47.5 °/2-5 min	Paul & Chen '99
Broccoli	decreased yellowing	HWT	50 °/2 min	Forney '95
			45 °/10 min; 47 °/7.5 min	Tian et al. '96, '97
Collard	decreased yellowing	HAT	45 °/30 min	Wang '98
Green onions	inhibited elongation	HWT	55 °/2 min	Hong et al. '00
Guava	decreased softening and yellowing	HWT	46 °/35 min	McGuire '97
Kale	decreased yellowing	HAT	40 °/60 min	Wang '98
Potato	inhibited sprouting	HWT		Rangann et al. '98